



The only injectable
neostigmine in the market



Rumintral

Neostigmine in injectable solution



Product of choice in tympanism and gastrointestinal atony

- ✓ Treatment of ruminal overload, gastrointestinal atony and symptoms of meteorism
- ✓ Has effects on the respiratory system, increasing ciliary activity and bronchial secretions


Indispensable in the veterinary kit

- ✓ Maximum efficacy in treatments combined with Rumilab®

 **LABIANA**
always works



Mechanism of action of neostigmine



COMPOSITION PER ML

Neostigmine methylsulfate..... 1.5 mg

INDICATIONS

Bovines:

- Ruminal atony.
- Intestinal atony.

DOSAGE AND ROUTE OF ADMINISTRATION

0.22 mg/kg l.w. (equivalent to 0.15 ml/10 kg l.w. of Rumintral) by subcutaneous or intramuscular route.

WITHDRAWAL PERIOD

Meat: Zero days.

Milk: Zero days.

USE IN PREGNANCY AND LACTATION

Do not administer to females in pregnancy or in lactation.

CONTRAINDICATIONS

- Do not administer in cases of mechanical obstruction of the gastrointestinal or urinary tract.
- Do not administer jointly with other cholinesterase inhibitors, or with depolarizing neuromuscular blockers (succinylcholine).

SECONDARY EFFECT

Those inherent to cholinergic hypoactivity.

OVERDOSING

Marked muscular weakness, vomiting, colic, diarrhea, myosis, dyspnea, bradycardia, hypotension. Death is caused due to respiratory failure.

Antidote: Atropine.

SPECIAL PRECAUTIONS FOR STORAGE

Store in a cool, dry place and protected from light.

PRESENTATION

25 ml vial.

Registry no. 891 ESP

Medication subject to veterinary prescription.

Administration under veterinary control or supervision.